

The Erdős-Falconer distance problem on the unit sphere in vector spaces over finite fields

Le Anh Vinh
 Mathematics Department
 Harvard University
 Cambridge, MA 02138, US
 vinh@math.harvard.edu

October 9, 2008

Abstract

Hart, Iosevich, Koh and Rudnev (2007) show, using Fourier analysis method, that the finite Erdős-Falconer distance conjecture holds for subsets of the unit sphere in $\mathbb{F}_q^d$. In this note, we give a graph theoretic proof of this result.

1 Introduction

The Erdős Distance Problem is perhaps the best known problem in combinatorial geometry. How many distinct distances can occur among n points in the plane? Although this problem has received considerable attention, we are still far from the solution. The Falconer distance conjecture says that if $E \subset \mathbb{R}^d$, $d \geq 2$, has Hausdorff dimension greater than $\frac{d}{2}$, then the set of distances occur in E has positive Lebesgue measure. See [9] for the connections between the Erdős and Falconer distance conjectures.

In the finite field setting, the distance problem turns out to have features of both the Erdős and Falconer distance problems in real spaces. Let $\mathbb{F}_q$ denote the finite field with q elements where $q \gg 1$ is an odd prime power. For any $x, y \in \mathbb{F}_q^d$, the distance between x, y is defined as $\|x - y\| = (x_1 - y_1)^2 + \dots + (x_d - y_d)^2$. Let $E \subset \mathbb{F}_q^d$, $d \geq 2$. Then the finite analog of the classical Erdős distance problem is to determine the smallest possible cardinality of the set

$$\Delta(E) = \{\|x - y\| : x, y \in E\}, \quad (1.1)$$

viewed as a subset of $\mathbb{F}_q$. Bourgain, Katz and Tao ([4]), showed, using intricate incidence geometry, that for every $\varepsilon > 0$, there exists $\delta > 0$, such that if $E \subset \mathbb{F}_q^2$ and $C_\varepsilon^1 q^\varepsilon \leq |E| \leq C_\varepsilon^2 q^{2-\varepsilon}$, then $|\Delta(E)| \geq C_\delta |E|^{\frac{1}{2}+\delta}$ for some constants $C_\varepsilon^1, C_\varepsilon^2$ and C_δ . The relationship between ε and δ in their argument is difficult to determine. Going up to higher dimension

using arguments of Bourgain, Katz and Tao is quite subtle. Iosevich and Rudnev [8] establish the following result using Fourier analytic method.

Theorem 1.1 ([8]) *Let $E \subset \mathbb{F}_q^d$ such that $|E| \gtrsim Cq^{d/2}$ for C sufficiently large. Then*

$$|\Delta(E)| \gtrsim \min \left\{ q, \frac{|E|}{q^{(d-1)/2}} \right\}. \quad (1.2)$$

In view of this result, Iosevich and Rudnev ([8]) formulated the Erdős-Falconer conjecture as follows.

Conjecture 1.2 *Let $E \subset \mathbb{F}_q^d$ such that $|E| \geq C_\epsilon q^{\frac{d}{2}+\epsilon}$. Then there exists $c > 0$ such that $|\Delta(E)| \geq cq$.*

By modifying the proof of Theorem 1.1 slightly, Iosevich and Rudnev ([8]) obtain the following stronger conclusion.

Theorem 1.3 ([8]) *Let $E \subset \mathbb{F}_q^d$ such that $|E| \geq Cq^{\frac{d+1}{2}}$ for sufficient large constant C . Then $\Delta(E) = \mathbb{F}_q$.*

In [7], the authors show that Theorem 1.3 is essentially sharp, which implies that Conjecture 1.2 is not true in general. They show however, that the exponent predicted by Conjecture 1.2 does hold for subsets of the sphere $S^{d-1} = \{x \in \mathbb{F}_q^d : x_1^2 + \dots + x_d^2 = 1\}$.

Theorem 1.4 ([7]) *Let $E \subset \mathbb{F}_q^d$, $d \geq 3$, be a subset of the sphere*

$$S^{d-1} = \{x \in \mathbb{F}_q^d : \|x\| = 1\}.$$

Suppose that $|E| \geq Cq^{d/2}$ with a sufficiently large constant C . Then there exists $c > 0$ such that $|\Delta(E)| > cq$.

In this note, we will give a graph theoretic proof of this result. The rest of this note is organized as follows. In Section 2, we construct our main tools to study the Erdős-Falconer distance problem over subsets of the sphere, the graphs associated to the projective spaces over finite fields. Our construction follows one of Bannai, Shimabukuro and Tanaka in [3]. We then prove Theorem 1.4 in Section 3. We also call the reader's attention that this note is a subsequent of an earlier paper [11].

Remark 1.5 *If d is even, Hart, Iosevich, Koh and Rudnev showed that all the distances can be obtained under the same assumption and the size condition on E cannot be relaxed. If d is odd then we cannot in general get all the distances if $|E| \ll q^{\frac{d+1}{2}}$. Interested readers can see [7] for a detailed discussion and related results.*

2 Finite non-Euclidean graphs

In this section, we give a construction of graphs from the action of simple orthogonal group on the set of non-isotropic square-type of projective spaces over finite fields. Our construction follows one of Bannai, Shimabukuro and Tanaka in [3]. Let $V = \mathbb{F}_q^d$ be the d -dimensional vector space over the finite field $\mathbb{F}_q$ (q is an odd prime power). For each element x of V , we denote the 1-dimensional subspace containing x by $[x]$. Let Ω be the set of all square type non-isotropic 1-dimensional subspaces of V with respect to the quadratic form $Q(x) = x_1^2 + \dots + x_d^2$. The simple orthogonal group $O_d(\mathbb{F}_q)$ acts transitively on Ω , and yields a symmetric association scheme $\Psi(O_d(\mathbb{F}_q), \Omega)$ of class $(q+1)/2$. We have two cases.

Case I. Suppose that $d = 2m + 1$. The relations of $\Psi(O_{2m+1}(\mathbb{F}_q), \Omega)$ are given by

$$\begin{aligned} R_1 &= \{([U], [V]) \in \Omega \times \Omega \mid (U + V) \cdot (U + V) = 0\}, \\ R_i &= \{([U], [V]) \in \Omega \times \Omega \mid (U + V) \cdot (U + V) = 2 + 2\nu^{-(i-1)}\} \quad (2 \leq i \leq (q-1)/2) \\ R_{(q+1)/2} &= \{([U], [V]) \in \Omega \times \Omega \mid (U + V) \cdot (U + V) = 2\}, \end{aligned}$$

where ν is a generator of the field $\mathbb{F}_q$ and we assume $U \cdot U = 1$ for all $[U] \in \Omega$ (see [2]).

Case II. Suppose that $d = 2m$. The relations of $\Psi(O_{2m}(\mathbb{F}_q), \Omega)$ are given by

$$\begin{aligned} R_i &= \{([U], [V]) \in \Omega \times \Omega \mid (U + V) \cdot (U + V) = 2 + 2^{-1}\nu^i\} \quad (1 \leq i \leq (q-1)/2) \\ R_{(q+1)/2} &= \{([U], [V]) \in \Omega \times \Omega \mid (U + V) \cdot (U + V) = 2\}, \end{aligned}$$

where ν is a generator of the field $\mathbb{F}_q$ and we assume $U \cdot U = 1$ for all $[U] \in \Omega$ (see [2]).

The graphs (Ω, R_i) are not Ramanujan in general, but fortunately, they are asymptotic Ramanujan for large q . The following theorem summarizes the results from [3] in a rough form.

Theorem 2.1 ([3]) *The graphs (Ω, R_i) ($1 \leq i \leq (q+1)/2$) are regular of order $q^{d-1}(1 + o(1))/2$ and valency $Kq^{d-2}(1 + o(1))$. Let λ be any eigenvalue of the graph (Ω, R_i) with $\lambda \neq$ valency of the graph then*

$$|\lambda| \leq k(1 + o(1))q^{(d-2)/2},$$

for some $k, K > 0$ (In fact, we can show that $k = 2$ and $K = 1$ or $1/2$).

3 Graph theoretic proof of Theorem 1.4

Let E be a subset of the unit sphere $S^{d-1} = \{x \in \mathbb{F}_q^d : \|x\| = 1\}$ with $|E| \geq Cq^{d/2}$. Let $E_1 = \{[x] : x \in E\} \subset \Omega$ (where Ω is the set of all square type non-isotropic 1-dimensional subspaces of V with respect to the quadratic form $Q(x) = x_1^2 + \dots + x_d^2$). Since each line through origin in $\mathbb{F}_q^d$ intersects the unit sphere S^{d-1} at two points, $|E_1| \geq Cq^{d/2}/2$. Suppose that $([U], [V]) \in E_1 \times E_1$ is an edge of (Ω, R_i) . Then

$$(U + V) \cdot (U + V) = 2 + \alpha_i,$$

where $\alpha_i = 2\nu^{-(i-1)}$ if d is odd and $\alpha_i = 2^{-1}\nu^i$ if d is even. Since $U \cdot U = V \cdot V = 1$, we have $(U - V) \cdot (U - V) = 2 - \alpha_i$. The distance between U and V (in E) is either $(U + V) \cdot (U + V)$ or $(U - V) \cdot (U + V)$, so

$$|\Delta(E) \cap \{2 + \alpha_i, 2 - \alpha_i\}| \geq 1.$$

Therefore, it is sufficient to show that $E_1 \times E_1$ contains edges of at least cq graphs among (Ω, R_i) , $1 \leq i \leq (q+1)/2$.

To complete the proof, we need the following result from spectral graph theory. We call a graph G (n, d, λ) -regular if G is a d -regular graph on n vertices with the absolute value of each of its eigenvalues but the largest one is at most λ . It is well-known that if $\lambda \ll d$ then a (n, d, λ) -regular graph behaves similarly as a random graph $G_{n,d/n}$. Presicely, we have the following result (see Corollary 9.2.5 and Corollary 9.2.6 in [1]).

Theorem 3.1 ([1]) *Let G be a (n, d, λ) -regular graph. For every set of vertices B of G , we have*

$$|e_G(B) - \frac{d}{2n}|B|^2| \leq \frac{1}{2}\lambda|B|, \quad (3.1)$$

where $e_G(B)$ is number of edges in the induced subgraph of G on B .

From Theorem 3.1, we have

$$|e_{(\Omega, R_i)}(E_1) - \frac{Kq^{d-2}(1+o(1))}{q^{d-1}(1+o(1))/2}|E_1|^2| \leq \frac{1}{2}k(1+o(1))q^{(d-2)/2}|E_1|.$$

Since $|E_1| \geq Cq^{d/2}/2$ for C sufficiently large, the left hand term $\frac{1}{2}k(1+o(1))q^{(d-2)/2}|E_1|$ is neglected by $\frac{Kq^{d-2}(1+o(1))}{q^{d-1}(1+o(1))/2}|E_1|^2$. Thus, we have $e_{(\Omega, R_i)}(E_1) = O(|E_1|^2/q)$. Besides $E_1 \times E_1$ is edge-decomposed into $(q+1)/2$ graphs (Ω, R_i) , $1 \leq i \leq (q+1)/2$. This implies that $E_1 \times E_1$ contains edges of at least cq graphs among (Ω, R_i) , $1 \leq i \leq (q+1)/2$ for some constant $c > 0$. The theorem follows.

References

- [1] N. Alon and J. H. Spencer, *The probabilistic method*, 2nd ed., Willey-Interscience, 2000.
- [2] E. Bannai, S. Hao and S.-Y. Song, Character tables of the association schemes of finite orthogonal groups acting on the nonisotropic points, *Journal of Combinatorial Theory, Series A* **54** (1990), 164-170.
- [3] E. Bannai, O. Shimabukuro and H. Tanaka, Finite analogues of non-Euclidean spaces and Ramanujan graphs, *European Journal of Combinatorics* **25** (2004), 243-259.
- [4] J. Bourgain, N. Katz, T. Tao, A sum-product estimate in finite fields, and applications, *Geom. Funct. Anal.* **14** (2004), 27-57.

- [5] P. Brass, W. Moser and J. Pach, *Research problems in discrete geometry*, Springer, 2005.
- [6] P. Erdős, On sets of distances of n points, *Amer. Math. Monthly* **53** (1946) 248–250.
- [7] D. Hart, A. Iosevich, D. Koh and M. Rudnev, Averages over hyperplanes, sum-product theory in vector spaces over finite fields and the Erdős-Falconer distance conjecture, preprint, 2007.
- [8] A. Iosevich, M. Rudnev, Erdős distance problem in vector spaces over finite fields, *Transactions of the American Mathematical Society* **359** (12) (2007), 6127-6142.
- [9] A. Iosevich, M. Rudnev and I. Uriarte-Tuero. Theory of dimension for large discrete sets and applications, preprint, 2007.
- [10] L. A. Székely, Crossing numbers and hard Erdős problems in discrete geometry, *Comb. Probab. Comput.* **6** (1997), 353–358.
- [11] L. A. Vinh, Explicit Ramsey graphs and Erdős distance problem over finite Euclidean and non-Euclidean spaces, *Electronic Journal of Combinatorics* **15** (2008), R5.